



# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending December 9, 2017

Prepared by the Infectious Disease Epidemiology and Outbreak Response Bureau  
Prevention and Health Promotion Administration  
Maryland Department of Health

*The data presented in this document are provisional and subject to change as additional reports are received.*

## SUMMARY

During the week ending December 9, 2017, influenza-like illness (ILI) intensity in Maryland was **MINIMAL** and there was **REGIONAL** geographic activity. The proportion of outpatient visits for ILI reported by Sentinel Providers and outpatient visits for ILI at Maryland Emergency Departments both increased slightly. The proportion of MRITS respondents reporting ILI was low. Clinical laboratories reported an increase in the proportion of specimens testing positive for influenza. Thirty-four specimens tested positive for influenza at the MDH lab. There were 35 influenza-associated hospitalizations. One respiratory outbreak was reported to MDH.


[Click here to visit our influenza surveillance web page](#)

ILI Intensity Levels
✓ Minimal
Low
Moderate
High

Influenza Geographic Activity
No Activity
Sporadic
Local
✓ Regional
Widespread

## ILINet Sentinel Providers


Twenty four sentinel providers reported a total of 7,420 visits this week. Of those, 118 (1.6%) were visits for ILI. This is **below** the Maryland baseline of **2.0%**.



ILI Visits To Sentinel Providers By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	31 (26%)	36 (33%)	285 (31%)
Age 5-24	54 (46%)	42 (39%)	351 (38%)
Age 25-49	21 (18%)	21 (19%)	180 (19%)
Age 50-64	8 (7%)	6 (6%)	76 (8%)
Age ≥ 65	4 (3%)	3 (3%)	38 (4%)
<b>Total</b>	<b>118 (100%)</b>	<b>108 (100%)</b>	<b>930 (100%)</b>

## Visits to Emergency Departments for ILI

Emergency Departments in Maryland reported a total of 44,041 visits this week through the [ESSENCE surveillance system](#). Of those, 961 (2.2%) were visits for ILI.



ILI Visits To Emergency Departments By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	371 (39%)	322 (35%)	2,626 (36%)
Age 5-24	204 (21%)	233 (26%)	1,851 (25%)
Age 25-49	210 (22%)	204 (22%)	1,705 (23%)
Age 50-64	94 (10%)	92 (10%)	714 (10%)
Age ≥ 65	82 (9%)	57 (6%)	495 (7%)
<b>Total</b>	<b>961 (100%)</b>	<b>908 (100%)</b>	<b>7,391 (100%)</b>

### Neighboring states' influenza information:


- Delaware <http://dhss.delaware.gov/dph/epi/influenzahome.html>
- District of Columbia <http://doh.dc.gov/service/influenza>
- Pennsylvania <http://www.health.pa.gov/My%20Health/Diseases%20and%20Conditions/I-L/Pages/Influenza.aspx#.V-LtaPkrJD8>
- Virginia <http://www.vdh.virginia.gov/epidemiology/influenza-flu-in-virginia/influenza-surveillance/>
- West Virginia <http://dhhr.wv.gov/oeps/disease/flu/Pages/fluSurveillance.aspx>

# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending December 9, 2017

## Community-based Influenza Surveillance (MRITS)


MRITS is the Maryland Resident Influenza Tracking System, a weekly survey for influenza-like illness (ILI). A total of 546 residents responded to the [MRITS survey](#) this week. Of those, 4 (0.7%) reported having ILI and missing greater than 7 cumulative days of regular daily activities.




MRITS Respondents Reporting ILI By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	--	--	3 (7%)
Age 5-24	3 (75%)	1 (25%)	13 (32%)
Age 25-49	--	--	5 (12%)
Age 50-64	1 (25%)	2 (50%)	15 (37%)
Age ≥ 65	--	1 (25%)	5 (12%)
<b>Total</b>	<b>4 (100%)</b>	<b>4 (100%)</b>	<b>41 (100%)</b>

## Clinical Laboratory Influenza Testing

There were 47 clinical laboratories reporting 1,814 influenza diagnostic tests, mostly rapid influenza diagnostic tests (RIDTs). Of those, 150 (8.3%) were positive for influenza. Of those testing positive, 121 (81%) were influenza Type A and 29 (19%) were influenza Type B. The [reliability of RIDTs](#) depends largely on the conditions under which they are used. False-positive (and true-negative) results are more likely to occur when the disease prevalence in the community is low, which is generally at the beginning and end of the influenza season and during the summer.




Positive Rapid Flu Tests by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A	121 (81%)	78 (74%)	338 (68%)
Type B	29 (19%)	28 (26%)	160 (32%)
<b>Total</b>	<b>150 (100%)</b>	<b>106 (100%)</b>	<b>498 (100%)</b>



## State Laboratories Administration Influenza Testing

The MDH Laboratories Administration performed a total of 157 PCR tests for influenza and 34 (21.7%) were positive for influenza. Of those testing positive, 29 (85.3%) were positive for Type A (H3), 4 (11.8%) was positive for Type A (H1), and 1 (2.9%) was positive for Type B (Yamagata). PCR testing is more reliable than RIDT. The MDH testing identifies subtypes of influenza A and lineages of influenza B, information that is not available from the RIDT results. The table below summarizes results by type, subtype, and lineage.



Positive PCR Tests by Type (Subtype)	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A (H1)	4 (12%)	1 (7%)	8 (10%)
Type A (H3)	29 (85%)	10 (67%)	62 (78%)
Type B (Victoria)	--	--	--
Type B (Yamagata)	1 (3%)	4 (27%)	7 (9%)
Type A (H3N2v)	--	--	3 (4%)
<b>Total</b>	<b>34 (100%)</b>	<b>15 (100%)</b>	<b>80 (100%)</b>

## Where to get an influenza vaccination


Interested in getting a flu vaccine for the 2017-18 influenza season? Go to <https://phpa.health.maryland.gov/influenza/Pages/getvaccinated.aspx> and click on your county/city of residence. You will be redirected to your local health department website for local information on where to get your flu vaccine.

# Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending December 9, 2017

## Influenza-associated Hospitalizations


A total of 35 influenza-associated hospitalizations were reported this week. (A person with an overnight hospital stay along with a positive influenza test of any kind, e.g., RIDT or PCR, is considered an “influenza-associated hospitalization” for purposes of influenza surveillance.)



Influenza-Associated Hospitalizations by Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	3 (9%)	4 (13%)	15 (10%)
Age 5-17	--	--	5 (3%)
Age 18-24	1 (3%)	--	5 (3%)
Age 25-49	6 (17%)	8 (25%)	26 (17%)
Age 50-64	6 (17%)	4 (13%)	31 (20%)
Age ≥ 65	19 (54%)	16 (50%)	70 (46%)
<b>Total</b>	<b>35 (100%)</b>	<b>32 (100%)</b>	<b>152 (100%)</b>

## Outbreaks of Respiratory Disease

There was one respiratory outbreak reported to MDH this week. (Disease outbreaks of any kind are reportable in Maryland. Respiratory outbreaks may be reclassified once a causative agent is detected, e.g., from ILI to influenza.)



Respiratory Outbreaks by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Influenza	1 (100%)	1 (100%)	4 (36%)
Influenza-like Illness	--	--	3 (27%)
Pneumonia	--	--	4 (36%)
Other Respiratory	--	--	--
<b>Total</b>	<b>1 (100%)</b>	<b>1 (100%)</b>	<b>11 (100%)</b>

## National Influenza Surveillance (CDC)

During week 49 (December 3-9, 2017), influenza activity increased in the United States.

- **Viral Surveillance:** The most frequently identified influenza virus type reported by public health laboratories during week 49 was influenza A. The percentage of respiratory specimens testing positive for influenza in clinical laboratories increased.
- **Novel Influenza A Virus:** One human infection with a novel influenza A virus was reported.
- **Pneumonia and Influenza Mortality:** The proportion of deaths attributed to pneumonia and influenza (P&I) was below the system-specific epidemic threshold in the National Center for Health Statistics (NCHS) Mortality Surveillance System.
- **Influenza-associated Pediatric Deaths:** One influenza-associated pediatric death was reported.
- **Influenza-associated Hospitalizations:** A cumulative rate of 4.3 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported.
- **Outpatient Illness Surveillance:** The proportion of outpatient visits for influenza-like illness (ILI) was 2.7%, which is above the national baseline of 2.2%. Seven of the 10 regions reported ILI at or above region-specific baseline levels. Four states experienced high ILI activity; five states experienced moderate ILI activity; New York City, Puerto Rico, and 16 states experienced low ILI activity; 25 states experienced minimal ILI activity; and the District of Columbia had insufficient data.
- **Geographic Spread of Influenza:** The geographic spread of influenza in 12 states was reported as widespread; Puerto Rico and 26 states reported regional activity; 10 states reported local activity; the District of Columbia, the U.S. Virgin Islands and two states reported sporadic activity; and Guam did not report.

